

## Akt (pan) Rabbit mAb (AR1599)

### Key Features

Host Species:	Rabbit
Reactivity:	Human, Mouse, Rat
Applications:	WB, IHC, IF, IP, ELISA
Isotype:	IgG, Kappa
MW:	55kD (Calculated) 55kD (Observed)

### Recommended Dilution Ratios

IHC:	1:200-1000
WB:	1:2000-10000
IF:	1:200-1000
ELISA:	1:5000-20000
IP:	1:50-200

### Storage

-15°C to -25°C/1 year (Do not lower than -25°C)

### Basic Information

Clonality	Monoclonal
-----------	------------

### Immunogen Information

Specificity	Endogenous
-------------	------------

### Target Information

Gene name	AKT1/AKT2/AKT3 RAC-alpha serine/threonine-protein kinase; RAC-beta serine/threonine-protein kinase; RAC gamma serine/threonine-protein kinase
Protein Name	

Organism	Gene ID	UniProt ID
Human	207; 208; 10000	P31749; P31751; Q9Y243
Mouse	11651; 11652; 23797	P31750
Rat	24185; 25233; 29414	P47196; P47197; Q63484

## Cellular Localization


Cytoplasm . Nucleus . Cell membrane . Nucleus after activation by integrin-linked protein kinase 1 (ILK1). Nuclear translocation is enhanced by interaction with TCL1A.

Phosphorylation on Tyr-176 by TNK2 results in its localization to the cell membrane where it is targeted for further phosphorylations on Thr-308 and Ser-473 leading to its activation and the activated form translocates to the nucleus. Colocalizes with WDFY2 in intracellular vesicles (PubMed:16792529).

Expressed in prostate cancer and levels increase from the normal to the malignant state (at protein level). Expressed in all human cell types so far analyzed. The Tyr-176 phosphorylated form shows a significant increase in expression in breast cancers during the progressive stages i.e. normal to hyperplasia (ADH), ductal carcinoma in situ (DCIS), invasive ductal carcinoma (IDC) and lymph node metastatic (LNMM) stages

## Tissue specificity

## Validation Data



Various whole cell lysates were separated by 4-20% SDS-PAGE, and the membrane was blotted with anti-AKT antibody. The HRP-conjugated Goat anti-Rabbit IgG(H + L) antibody was used to detect the antibody.

Lane 1: HeLa

Lane 2: Jurkat

Lane 3: C6

Lane 4: Mouse brain

Predicted band size: 55kDa

Observed band size: 55kDa

Immunofluorescence analysis of human-stomach tissue.

1, Akt Monoclonal Antibody(red) was diluted at

1:200(4°C, overnight).

2, Cy3 labeled Secondary antibody was diluted at 1:300(room temperature, 50min).

3, Picture B: DAPI(blue) 10min.

Picture A: Target. Picture B: DAPI. Picture C: merge of A+B

Immunofluorescence analysis of rat-lung tissue.



1, Akt Monoclonal Antibody(red) was diluted at

1:200(4°C, overnight).

2, Cy3 labeled Secondary antibody was diluted at 1:300(room temperature, 50min).

3, Picture B: DAPI(blue) 10min.

Picture A: Target. Picture B: DAPI. Picture C: merge of A+B



Immunofluorescence analysis of mouse-spleen tissue.

1, Akt Monoclonal Antibody (red) was diluted at 1:200 (4°C, overnight).

2, Cy3 labeled Secondary antibody was diluted at 1:300 (room temperature, 50 min).

3, Picture B: DAPI (blue) 10 min.



Picture A: Target. Picture B: DAPI. Picture C: merge of A+B

Immunofluorescence analysis of HEK293.

Picture A: AKT antibody (red).

Picture B: DAPI (blue).

Picture C: Merge of A+B



For Research Use Only